

Effects of Monacolin K of Red Rice and Glucomannan, Combined with a Low Calorie Diet, in Treatment of Dyslipidemia and Hypertension

Teresa Esposito, Salvatore Allocca, Laura Adelfi, Giovanni Messina, Marcellino Monda* and Bruno Varriale

Department of Experimental Medicine, Second University of Naples, Naples, Italy

Abstract

Alarming increase in incidence of cardiovascular disease is result of a nutrition pattern characterized by an increase in consumption of fats, cholesterol, sugar and other refined carbohydrates, concomitant with low consumption of polyunsaturated fatty acids and fibers. The purpose of this study was to evaluate the efficacy of the monacolin K of red rice and glucomannan, associated with low-calorie diet, on improvement of lipid profile and hypertension, symptoms of metabolic syndrome. In 180 Neapolitan patients with metabolic syndrome, we evaluated effects of monacolin K of red rice and glucomannan, associated with a low-calorie diet, on total cholesterol (CT), LDL cholesterol (LDL), HDL cholesterol (HDL), triglycerides (TG) and blood pressure (BP). The results showed a significant decrease in average value of CT (-20%), LDL (-25%), TG (-20%), with an increase in HDL (+15%) and a decrease in BP (-20%). These findings corroborate evidences showing a therapeutic effects of these nutraceuticals in the treatment of dyslipidemia and hypertension, when a low-calorie diet alone is ineffective.

Keywords: Hypertension; Dyslipidemia; Cholesterol; Blood pressure

Introduction

Metabolic syndrome is a medical condition deserving of special attention because of its prevalence and impact [1]. This term does not indicate a single disease, but a group of predisposing factors which, joined together, place the subject in a band of high risk for diseases such as diabetes, cardiovascular problems and hepatic steatosis [2-5].

Numerous epidemiological and experimental studies have demonstrated that metabolic syndrome affects almost half of adults over 50-60 years [4,6-12]. Probably, this incidence will grow in the future, considering spreading childhood obesity. An excess of body fat, especially when it is localized in the abdominal region, leads to an imbalance of the metabolism of fats and sugars that induces hyperinsulinemia (high insulin level in the blood, an indicator of increased resistance to this hormone) [9,13]. In addition to strategies based on changing lifestyle, it is frequent utilization of natural substances, with beneficial properties [14-18]. Monacolin K of red rice (10 mg, 90% by policosanol) is able to inhibit the HMG-CoA reductase, which is a key enzyme in the biosynthesis of cholesterol. Glucomannan (3 g per day) is particularly useful in the reduction of body weight and cholesterol [19]. It is also useful in constipation to regulate bowel function alternately in irritable bowel syndrome. The use of these products called "nutraceuticals" would appear useful in patients with poor tolerance to traditional drugs or those who reject traditional drugs [20-23]. The purpose of the study was to evaluate efficacy of the monacolin K of red rice and glucomannan, associated with low-calorie diet, on improvement of lipid profile and hypertension, symptoms of metabolic syndrome.

Materials and Methods

180 subjects (83 females and 97 males) with mild to moderate hypercholesterolemia, (range of total cholesterol between 200 and 290 mg/dl), mild hypertension (no antihypertensive drug therapy; 140-159/90-99 mm Hg systolic/diastolic blood pressure), in which low-fat diet for three months did not achieve the treatment goal, were treated also with monacolin K of red rice (10 mg, 90% in policosanol per day)

and fiber glucomannan (3 g per day) for 24 weeks. Blood pressure (BP), lipid profile [total cholesterol (CT), LDL cholesterol (LDL), HDL cholesterol (HDL), triglycerides (TG)], biochemical parameters related to tolerability (GOT, GPT, CPK) and body mass index were determined at baseline (T0), to 12th week (T1) and 24th week (T2). All biochemical parameters were performed by a single laboratory accredited to ISO parameters accuracy. All subjects expressed informed consent. The data were expressed as mean \pm standard deviation (M \pm DS) and percentage (%) values. Significance of differences between groups was determined by Student's test for paired data, for linear data and the X2 for nonparametric data (Table 1).

Results

The results are reported in table. Monacolin K of red rice and fiber glucomannan, associated with a low-calorie diet, for 24 weeks showed a reduction in CT, LDL and systolic-diastolic BP. At T1 COL was reduced by 12.8% in males and 13% females, LDL was reduced by 15% in males and 16.2% females. HDL was increased by 6.1% in males and 6.6% females, TG were reduced by 14.9% in males and 13% females. BP was decreased by 20% for both diastolic and systolic values. At T2 COL was reduced by 19.3% in males and 19.5% in females, LDL was reduced by 25.2% in males and 27% females; HDL was increased by 11.2% in males and 15.4% females. TG were reduced by 22% in males and 21% females. Body weight decreased in association to decrease in lipid parameters.

***Corresponding author:** Marcellino Monda, Professor, Department of Experimental Medicine, Section of Human Physiology, and Clinical Dietetic Service, Second University of Naples, Via Costantinopoli 16, 80138 Naples, Italy, Tel: +39 +81 566 5804; Fax: +39 +81 5665841; E-mail: marcellino.monda@unina2.it

Received July 27, 2015; **Accepted** September 05, 2015; **Published** September 12, 2015

Citation: Esposito T, Allocca S, Adelfi L, Messina G, Monda M, et al. (2015) Effects of Monacolin K of Red Rice and Glucomannan, Combined with a Low Calorie Diet, in Treatment of Dyslipidemia and Hypertension. Biol Med (Aligarh) 7: 250. doi: 10.4172/0974-8369.1000250

Copyright: © 2015 Esposito T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

	Basal	12th week		24th week		Statistical significance
	M ± DS	M ± DS	D % vs Basale	M ± DS	D % vs Basale	p
Total cholesterol (mg/dl)	262 ± 35	228 ± 32	-12.7	213 ± 27	-19.5	≥ 0.01
LDL cholesterol (mg/dl)	179 ± 29	149 ± 28	-15.6	135 ± 24	-25.9	≥ 0.01
HDL cholesterol (mg/dl)	52 ± 12	54 ± 14	6.7	55 ± 13	13.4	n.s
Triglycerides (mg/dl)	175 ± 13	135 ± 46	-14	136 ± 35	- 20.2	n.s
Systolic blood pressure (mmHg)	150 ± 9	141 ± 10	-9	117 ± 5	-33	≥ 0.01
Diastolic blood pressure (mmHg)	94 ± 4	87 ± 5	-7	77 ± 6	-17	≥ 0.01
Body mass index	28.1 ± 2.2	26.1 ± 2.5	-6.4	24.9 ± 2.3	-11,3	≥ 0.01

Table 1: Average values (SD) at baseline at 12th week and 24th week in 180 patients.

Discussion

High COL is one of many factors that predispose to cardiovascular diseases. Some of these factors are modifiable (smoking cigarette, blood pressure, diabetes mellitus), while others are called non-modifiable (age, sex, family history and genetic factors). The influence of diet on COL is on average equal to 15%, although significant modifications of the contribution dietary can cause variations up to a ± 30%. Possible failure of the diet requires use of lipid-lowering agents. Medicines used in presence of hypercholesterolemia are statins (HMG-CoA reductase) and fibrates (most useful in presence of high triglycerides).

An alternative strategy to classical pharmacological intervention is use of nutraceuticals with a good tolerability profile and activity on lipid profile [24]. A good effect on cholesterol synthesis is obtained by policosanol. These depress expression of HMG-CoA-reductase in concentration-dependent manner, probably by receptor-mediated mechanisms that inhibit transcription for this enzyme [25,26]. Some studies showed that policosanol induces other effects on cardiovascular risk, similar to pleiotropic effects attributed to statins [27]. Red rice to 5% in monacolin K is a natural substance that is used for thousands of years in China to achieve desirable values of cholesterol. Glucomannan is a polysaccharide with high molecular weight, is extracted from tuber of *Amorphophallus konjac*, a plant used in Japanese cuisine as an agent gelatifer. This fiber has the ability to attract a lot of water, increasing its volume up to 60-100 times and giving rise to a soft gelatinous mass. This feature gives glucomannan a dual effect: it reduces sense of hunger and absorption of fats and sugars, which are trapped in soft, viscous mass that forms in the intestine [27-34].

The present study showed that monacolin K of red rice and glucomannan is useful in therapy of dyslipidemia and hypertension, when a low-calorie diet alone is ineffective, and it confirm effectiveness of these nutraceuticals. Since low-fat diet alone for three months (before integration of monacolin K and glicomannan) did not achieve effects on metabolic parameters, the improvement in the lipid results could be attributed to a direct effect of monacolin K and glucomannan. In perspective, this experiment should be extended to younger subjects before appearance of metabolic syndrome. In this way, we could assess possible positive effects in prevention of metabolic syndrome.

References

1. Grundy SM, Brewer Jr HB, Cleeman JI, Smith Jr SC, Lenfant C, American Heart Association, et al. (2004) Definition of metabolic syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association

- conference on scientific issues related to definition. *Circulation* 109: 433–438.
2. Messina G, Dalia C, Tafuri D, Monda V, Palmieri F, et al. (2014) Orexin-A controls sympathetic activity and eating behavior. *Front Psychol* 5: 997.
3. Messina G, Vicidomini C, Viggiano A, Tafuri D, Cozza V, et al. (2012) Enhanced parasympathetic activity of sportive women is paradoxically associated to enhanced resting energy expenditure. *Auton Neurosci Basic Clin* 169: 102–106.
4. Monda M, Messina G, Vicidomini C, Viggiano A, Mangoni C, et al. (2006) Activity of autonomic nervous system is related to body weight in pre-menopausal, but not in post-menopausal women. *Nutr Neurosci* 9: 141–145.
5. Viggiano A, Vicidomini C, Monda M, Carleo D, Carleo R, et al. (2009) Fast and low-cost analysis of heart rate variability reveals vegetative alterations in noncomplicated diabetic patients. *J Diabetes Complications* 23: 119–123.
6. Messina G, Viggiano A, De Luca V, Messina A, Chieffi S, et al. (2013) Hormonal changes in menopause and orexin-a action. *Obstet Gynecol Int* 209812.
7. Messina G, De Luca V, Viggiano A, Ascione A, Iannaccone T, et al. (2013) Autonomic nervous system in the control of energy balance and body weight: personal contributions. *Neurol Res Int* 1–5.
8. Viggiano A, Nicodemo U, Viggiano E, Messina G, Viggiano A, et al. (2010) Mastication overload causes an increase in O₂ - production into the subnucleus oralis of the spinal trigeminal nucleus. *Neuroscience* 166: 416–421.
9. Monda M, Messina G, Mangoni C, De Luca B (2008) Resting energy expenditure and fat-free mass do not decline during aging in severely obese women. *Clin Nutr* 27: 657–659.
10. Monda M, Messina G, Scognamiglio I, Lombardi A, Martin GA, et al. (2014) Short-Term Diet and Moderate Exercise in Young Overweight Men Modulate Cardiac and Hepatocarcinoma Survival by Oxidative Stress. *Oxid Med Cell Longev* 1–7.
11. Di Bernardo G, Messina G, Capasso S, Del Gaudio S, Cipollaro M, et al. (2014) Sera of overweight people promote in vitro adipocyte differentiation of bone marrow stromal cells. *Stem Cell Res Ther* 5: 4.
12. Esposito M, Serpe FP, Diletti G, Messina G, Scortichini G, et al. (2014) Serum levels of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans and polychlorinated biphenyls in a population living in the Naples area, southern Italy. *Chemosphere* 94: 62–69.
13. Viggiano A, Chieffi S, Tafuri D, Messina G, Monda M, De Luca B (2013) Laterality of a second player position affects lateral deviation of basketball shooting. *J Sports Sci* 37–41.
14. Chen CH, Yang JC, Uang YS, Lin CJ. (2013) Improved dissolution rate and oral bioavailability of lovastatin in red yeast rice products. *Int J Pharm* 444: 18-24.
15. Venero CV, Venero JV, Wortham DC, Thompson PD. (2010) Lipid-lowering efficacy of red yeast rice in a population intolerant to statins. *Am J Cardiol* 105: 664–666.
16. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, et al. (2004) Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 114: 1752–1761.

17. Keaney Jr JF, Larson MG, Vasan RS, Wilson PW, Lipinska I, et al. (2003) Obesity and systemic oxidative stress: clinical correlates of oxidative stress in the Framingham Study. *Arterioscler Thromb Vasc Biol* 23: 434–439.
18. Becker DJ, Gordon RY, Halbert SC, French B, Morris PB, et al. (2009) Red yeast rice for dyslipidemia in statin-intolerant patients: randomized trial. *Ann Intern Med* 150: 830-839.
19. EFSA. Scientific Opinion on the substantiation of health claims related to monacolin K from red yeast rice and maintenance of normal blood LDL cholesterol concentrations (ID 1648, 1700) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J.* 2011;9:2304.
20. De Luca V, Viggiano E, Messina G, Viggiano A, Borlido C, et al. (2008) Peripheral amino acid levels in schizophrenia and antipsychotic treatment. *Psychiatry Investig* 5: 203–208.
21. Monda M, Amaro S, Sullo A, De Luca B (1995) Injection of muscimol in the posterior hypothalamus reduces the PGE1-hyperthermia in the rat. *Brain Res Bull* 37: 575–580.
22. Monda M, Viggiano A, Viggiano A, Viggiano E, Messina G, et al. (2007) Sympathetic and hyperthermic reactions by orexin A: Role of cerebral catecholaminergic neurons. *Regul Pept* 139: 39–44.
23. Monda M, Viggiano A, Viggiano A, Viggiano E, Messina G, et al. (2006) Quetiapine lowers sympathetic and hyperthermic reactions due to cerebral injection of orexin A. *Neuropeptides* 40: 357–363.
24. Menendez R, Arruzazabala ML, Mas R et al. (1997) Cholesterol-lowering effect of policosanol on rabbits with hypercholesterolemia induced by a starch-casein diet. *Br J Nutr* 77: 923-32.
25. Arruzazabala ML, Carbajal D, Mas R, Del Río A, Amor AM, et al. (1994) Cholesterol-lowering effect of policosanol in rabbits. *Biol Res* 27: 205-208.
26. Heber D et al. (1999) Cholesterol-lowering effects of a proprietary Chinese red-yeast-rice dietary supplement. *Am J Clin Nutr* 69: 231-236.
27. Doi K (1990) Effect of Konjac fibre (glucomannan) on glucose and lipids. *Eur J Clin Nutr* 49: 190-197.
28. Blundell JE, Burley VJ (1987) Satiating, satiety and the action of fibre on food intake. *Int J Obes* 11: 9-25.
29. Monda M, Viggiano A, Viggiano A, Fuccio F, De Luca V (2004) Injection of orexin A into the diagonal band of Broca induces sympathetic and hyperthermic reactions. *Brain Res* 1018: 265–271.
30. Monda M, Amaro S, Sullo A, De Luca B (1994) Posterior hypothalamic activity and cortical control during the PGE1 hyperthermia. *Neuroreport* 6: 135–139.
31. Monda M, Pittman QJ (1993) Cortical spreading depression blocks prostaglandin E1 and endotoxin fever in rats. *Am J Physiol* 264: R456–R459.
32. Esposito T, Uccello R, Caliendo R, Di Martino GF, Gironi Carnevale UA, et al. (2009) Estrogen receptor polymorphism, estrogen content and idiopathic scoliosis in human: a possible genetic linkage. *J Steroid Biochem Mol Biol* 116: 56-60.
33. Esposito T, Varriale B, D'Angelo R, Amato A, Sidoti A (2014) Regulation of flavin-containing mono-oxygenase (Fmo3) gene expression by steroids in mice and humans. *Horm Mol Biol Clin Investig* 20: 99-109.
34. Esposito T, Napoleone A, Allocca S, Varriale B, Monda M (2014) Diet Therapy of Obesity: Observations on the Usefulness of Weekly Supervision in the Improvement of Weight Loss. *J Obes Weight Loss Ther* 4: 225.