

Original Paper

## GRAZING ACTIVITY OF THE SEA URCHIN *TRIPNEUSTES GRATILLA* IN TROPICAL SEAGRASS BEDS OF BUTON ISLAND, SOUTHEAST SULAWESI, INDONESIA

Ma'ruf Kasim\*

Faculty of Fishery and Marine Science, Haluoleo University, Indonesia

Received : September, 8<sup>th</sup> 2009 ; Accepted : September, 30<sup>th</sup> 2009

### ABSTRACT

The research on grazing activity of a sea urchin (*Tripneustes gratilla*) in tropical seagrass beds was conducted in Buton Island. The objectives of the research were to study the grazing activity such as food preference, grazing rate and grazing times of *T. gratilla* in tropical seagrass community. The results of the research showed that *Thalassia hemprichii* and *Enhalus acoroides* are primary food items of *T. gratilla* and contained on average of 55 % and 31 % of the gut contents, respectively. The grazing rate of an individual ranged from 9.6 - 14.1 g wet weight in 24 hours. The grazing times of *T. gratilla* were between 01.00 - 11.00 a.m. and 15.00 - 21.00 p.m.

**Key words :** Grazing, *Tripneustes gratilla*, seagrass, gut contents

**Correspondence :** Phone : + 62-85E-mail : [marufkasim@hotmail.com](mailto:marufkasim@hotmail.com)

### INTRODUCTION

Seagrass beds play an important role in tropical marine coastal ecosystems and its high productivity contributes to lives of many organisms (den Hartog 1970). Seagrasses serve as habitats for epiphyte and epifauna, and also for macrobenthic organisms such as the sea urchin *Tripneustes gratilla* as feeding ground, shelters, nursery ground, and/or spawning ground (Mukai and Nojima, 1985; Mukai, *et al.*, 1987, Vaitilingon, 2003).

*T. gratilla* is a macrobenthic animal using seagrass beds as grazing area. *T. gratilla* in seagrass beds grazes on seagrass leaves (Eklöfa, 2008, Vaitilingon, 2003, Aziz 1994, Mukai and Nojima, 1985; Mukai *et al.*, 1987). Herring (1972) reported the analysis of gut

contents of the sea urchin in Zanzibar. He showed that seagrass leaf was a main food of the sea urchin. In the seagrass *Thalassodendron ciliatum* meadows of Mombasa Lagoon (Kenya), 39 % of the seagrass cover was heavily grazed by *T. gratilla* and more than 70 % of seagrass was found as dead shoots (Alcoverro and Mariani, 2002). *T. gratilla* was heavily grazing on seagrass *Halophila stipulacea* in Sinai seagrass beds, northern Red Sea and the Jordanian coast of the Gulf Aqaba (Lipkin 1979; Bouchon 1980; Hulings and Kirkman, 1982; Jafari and Mahasneh 1984). In Bolinao seagrass beds, Philippines, however, *T. gratilla* chose the seagrass leaves as

alternative foods among many food items (Klumpp *et al.*, 1993). Others that also live and graze on the seagrass beds are *Lytechinus variegatus* (Moore and McPherson 1965; Camp *et al.*, 1973; Greenway 1976; Bach 1979; Vadas *et al.*, 1982; Keller 1983; Greenway 1995), *Astropyga magnifica* (Bak and Nojima 1980), *Paracentrotus lividus* (Kirkman and Young 1981), *Diadema antillarum* (Zieman *et al.*, 1984), *Temnopleuris michaelsonii* (Cambridge *et al.*, 1986), *Tripneustes ventricosus* (Keller 1983; Tertschnig 1989), *Salmacis sphaeroides* (Klumpp *et al.*, 1993), and *Heliocidaris erythrogramma* (Larkum and West 1990).

*T. gratilla* in some areas has been known as an alternative food for human and this has been exploited by local community. That is a reason for more and more reducing of the sea urchin population. Mukai *et al.* (1987) reported that local people in Loloata Island, Papua New Guinea, collected the sea urchin and other benthic animal occasionally, but the effect of this exploitation may not be substantial. In Lombok and Sumbawa Island, Indonesia, the local people collect the sea urchins that served as food alternatively (Darsono and Sukarno 1993). If the population density of *T. gratilla* at a certain seagrass habitat becomes extremely high, it would block the balance of seagrass communities (Bak and Nojima, 1980). Lawrence (1975) reported when *Tripneustes ventricosus* reaches high population in a seagrass bed, the grazing activity left many bare ground areas. The grazing activity by sea urchin is important in determining the impact of herbivores invertebrates on seagrass assemblages in many coastal ecosystems.

The present study was undertaken to investigate the grazing activity of *T. gratilla* including its main food, food preference, grazing ability, and grazing times.

## MATERIALS AND METHODS

### Study sites

The field research was conducted in the tropical seagrass beds of Buton Island (5° 28' N; 122° 37' E), southeast Sulawesi, Indonesia (**Fig. 1**). Further analysis of food items of seurchin was held in the of biology laboratory Haluoleo University, Kendari.

The seagrass bed of study site was dominated by *Enhalus acoroides* and *Thalassia hemprichii*, and *T. gratilla* was usually found on the bottom of the seagrass meadows or sometimes climbing on the *E. acoroides* leaves. Most of sea floor was covered by fine sand and little piece of dead corals. The range of salinity and temperature was 30 – 31 ‰ and 26 – 30 °C, respectively. The current speed was between 2.6 – 3.4 cm s<sup>-1</sup>, and tidal range was 171 cm in average (Kasim, 1999).

### Food preference

To determine food items, which were dominant in gut contents of *T. gratilla*, 20 sea urchins were collected from seagrass beds. The various sizes of sea urchin were separated on the two different samples. < 40 mm was categorized in to small size sea urchin, and > 40 mm was categorized in to large sized of sea urchin. The size range of sea urchin was measured in diameter size with calipers.

A dissection for the specimens was carried out very carefully, the gut contents of *T. gratilla* were taken out and preserved in 70 % alcohol. The gut contents were observed by using a stereomicroscope. Each food type was separated to measure its volume and wet weight. Food items in the gut contents were identified comparing with leaf and thallus tissues of all seagrass and sea weed species found in the field (Kasim, 1999). For analyzing gut contents, an index of preponderance (IP) (Natarajan and Jhingran,

1961) and an Ivlev's electivity index (IE) (Ivlev, 1961) were used.

$$IP = \frac{V_i \times O_i}{\sum (V_i \times O_i)} \times 100$$

Where  $V_i$  is composition ratio of  $i$ -th food type in volume,  $O_i$  is occurrence frequency of  $i$ -th food type. IE is ;

$$IE = \frac{r_i - p_i}{r_i + p_i}$$

Where  $r_i$  is expressed the volume percentage of  $i$ -th food type in gut contents,  $p_i$  is relative percentage of  $i$ -th food type in volume in the field (seagrass bed). The IE value can vary between  $-1$  and  $+1$ . The positive IE means bigger proportion in the gut contents than in the field, which shows a preference on that type of food.

### Grazing rates

In order to measure the grazing rate, each individual of *T. gratilla* was put in the cage that made from the metal and net (1 mm mesh size) with size 60 x 60 x 50 cm (length, width and height). Eight sea urchins and eight cages were prepared for the experiment. Eight cages were securely put on the bare sand bottom near the original seagrass bed. Eight individuals of *T. gratilla* were collected from seagrass beds. In order to determine the grazing rates related to the seaurchin sizes, eight samples were divided into two groups. Four seaurchin were categorizing in small size (< 2.5 cm diameter) and four sea urchin in large size (> 5.5 cm diameter).

A handful of fresh seagrass leaves (including three species, *E. acoroides*, *Th. hemprichii* and *C. serrulata*) were weighed in wet weight at start point ( $W_0$ ). Just before weighing, the selected seagrass leaves

(complete and young leaves) were cleaned from the periphyton and macroalga which attach on the leaves. A young complete leaf was selected to avoid some physical factor that possibility to washing or damage some small parts of leaves. These methods to decrease any error data (weight) after 24 hours. Then, the seagrass leaves were put in a cage with stones as sinker to serve as food for *T. gratilla*. Then it was put into cages, one for each cage. After 24 hours, seagrass leaves were collected and weighed to know the final seagrass weight ( $W_{24}$ ). The supplying and weighing of seagrass leaves were carried out every 24 hours intensively for 10 days to gain the grazing rates of *T. gratilla*. The formulation of the grazing rates,  $G$ , is :

$$G = W_0 - W_{24} \\ (\text{g seagrass (WW) ind}^{-1} \text{ day}^{-1})$$

As preliminary research in November 1998, the weight of some groups of seagrass leaves without *T. gratilla* in a cage were have no significant different after 24 hours (Kasim, 1999). Its means, no other factors were decrease the weigh of seagrass leaves aside the grazing of *T. gratilla* after 24 hours.

### Diurnal Grazing Activity

Diurnal grazing activity of *T. gratilla* was investigates by two cages, 100 x 100 x 50 cm (no net at bottom) and were put onto dense seagrass meadows. Twenty four individuals of *T. gratilla* (30 – 50 mm) were gathered from the seagrass bed and kept in the cages (each cage contained 12 individuals).

After 24 hours, every two hours, two *T. gratilla* were picked up from cages (each cage 1 individual), and materials in the gut were taken out by cutting the gut and stored in bottles (preserved by 70 % alcohol). The volume of gut contents was measured with cylinder glass (50 ml).

## Measurement of water conditions

The water temperature on the bottom at the experimental sites was measured with a mercury thermometer, and salinity with a salinometer. Current velocity was measured at two points directly near the cages, just above the substratum surface using a kite sink drift modification (Kasim, 1999).

## Statistical analysis

The grazing rates of sea urchin between diameter in size (small size and large size) were analyzed by one-way analysis of variance (ANOVA). Standard correlation analysis was addressed to statistically evaluate the correlation between tidal level and volume of gut, with Spearman correlation coefficients ( $P \leq 0.05$ ) (Oliveira, *et al.*, 1997).

# RESULTS AND DISCUSSION

## RESULTS

### Food habits

*Thalassia hemprichii* had the largest IP value about 55 % and *Enhalus acoroides* was the second, about 32 % (Fig. 2). This showed that *T. hemprichii* was a main food, which was followed by *E. acoroides*. As a whole, seagrasses including *Halodule uninervis* and *Cymodocea serrulata* predominated in the gut contents.

### Food preference

Some other foods were not seagrasses, such as algae and small animals. *T. hemprichii* had relative high index of electivity (IE), about 0.65. If it is compared to others species occurred in gut contents of *T. gratilla*, it can be considered that *T. hemprichii* leaf was preferred by *T. gratilla* (Table 1).

## Grazing rate

The grazing rates of *T. gratilla* varied, but tended to increase with size of *T. gratilla*. Small-sized sea urchins ( $n = 4$ ), 22.8 – 24.3 mm ( $m \pm sd$  :  $23.6 \pm 0.1$  mm) (A) and 21.9 – 23.6 mm ( $22.8 \pm 0.1$  mm) (B) in diameter, had average grazing rates of  $9.8 \pm 1.6$  and  $9.93 \pm 1.8$  g WW  $\text{ind}^{-1} \text{day}^{-1}$ , respectively. The large sized sea urchins ( $n = 4$ ), 51.9 – 53.8 mm ( $52.9 \pm 0.1$  mm) (C) and 54.8 – 56.2 mm ( $55.5 \pm 0.1$  mm) (D) in diameter, had grazing rates of  $13.8 \pm 2.7$  and  $14.3 \pm 2.4$  g WW  $\text{ind}^{-1} \text{day}^{-1}$ , respectively (Fig. 3).

The daily grazing rates of the sea urchins are not different in both similar diameter sizes. The small-sized and the large-sized *T. gratilla* had grazing rates between  $10.8 \pm 0.8$  (A),  $10.5 \pm 0.9$  (B),  $15.4 \pm 0.9$  (C), and  $15.5 \pm 1.0$  (D) g WW  $\text{ind}^{-1} \text{day}^{-1}$  (Fig. 4).

Analysis of variances of feeding rates on seagrasses showed no significant different between same diameter in size (ANOVA, AB,  $F=0.42$ ,  $P=0.533$ , CD,  $F=0.12$ ,  $P=0.739$ ; Tab.3). But for different diameter in size between small and large, there were significant different (ANOVA, AC,  $F=88.46$ ,  $P<0.001$ , BD,  $F=85.68$ ,  $P<0.001$ ; Tabel 2).

## Grazing time

Fig. 5 shows a diurnal fluctuation in volume of gut contents of *T. gratilla*. The fluctuation suggests that the grazing activity of *T. gratilla* had a tidal rhythm which corresponded to fluctuation of tidal level. The gut contents of *T. gratilla* decreased at high tides (11.00 a.m. – 13.00 a.m. and 23.00 to p.m.– 01.00 a.m). The speed of stream tended to increase or rise that made *T. gratilla* seen not eaten and heading on the under of dense seagrass leaves for preventing it from the strong stream. While at 01.00 a.m. – 09.00 a.m. and 15 p.m. – 21.00 p.m., the seawater surface started falling and also the speed of stream tended to decrease so this chance was used by *T. gratilla* more active to grazing at noon and evening,

particularly at 09.00 a.m.– 12 .00 a.m. and between 18.00 p.m. – 21.00 p.m.

To find out the correlation between tidal level and volume of gut content of *T. gratilla*, standard correlation analysis was performed on these data. From these analysis, the tendency shows negative correlation between tidal level and volume of gut sea urchin (correlation = -0.716), suggesting that the feeding activity of *T. gratilla* is negatively affected by the tidal level.

## DISCUSSION

The sea urchin, *T. gratilla*, showed a strong preference for seagrass leaves *Th. hemprichii* and *E. acoroides* than other macrophytes as indicated by the food habit and food preference of this sea urchin in tropical seagrass beds in Buton Island, Indonesia. *Th. hemprichii*, *E. acoroides*, *H. uninervis* and *C. serrulata*, were dominant seagrass species in this site. *T. gratilla* was the most common sea urchin in seagrass bed in Buton Island, and accounted for 80 % of the total number of sea urchins. The remaining 20 % addressed to *Diadema setosum*, *Echinometra mathei*, *Echinotrix diadema* and *salmacis* sp. (Kasim, 1999).

*T. gratilla* shows a strong preference for *Th. hemprichii* over *E. acoroides* or other seagrass such as *H. uninervis* and *C. serrulata*, as indicated by grazing activity on seagrass habitat. *T. gratilla* consistently consumes more *Th. hemprichii* throughout the 20 individuals experiment. Several studies have been reported to clarify the preferential foods of *T. gratilla*. The research carried out in Papua New Guinean seagrass beds by Hattori, *et al.*, (1985), was clarified that *T. gratilla* is an important element in tropical seagrass beds. It lives with grazing on seagrass leaves directly (Mukai, *et al.*, 1987). In Mombasa lagoon, 90 % of the total number species of sea urchin is *T. gratilla* that direct grazing on seagrass shoot and has a significant impact to the highest

number of seagrass dead shoot (Alcoverro and Mariani, 2002).

Lawrance (1975) reported that food preference pattern of the sea urchin was not clear but there was a tendency to favorite seagrasses, genus *Thalassia* and *Syringodium* in Atlantic seagrass beds. Mukai and Nojima (1985) reported that *Th. hemprichii* was a main food of *T. gratilla* in seagrass beds of Loloata Island, Papua New Guinea. From the field research in Zanzibar, Herring (1972) found about 82.5 % seagrass leaves in gut contents of sea urchin. Darsono and Sukarno (1993) reported that gut contents of *T. gratilla* in seagraas beds of Nusa Dua (Bali) were dominated by crushed and cut seagrass leaves, particularly *Th. hemprichii* and *E. acoroides*. They found a few algae and remain of *Ulva* sp. These foods preferential in seagrasses have been attribute to both food habit and food preference of *T. gratilla* in Buton Island seagrass beds. The occurrence and abundance of *Th. hemprichii* and *E. acoroides* in seagrass beds around the sea urchin has been forced to choose the kind of seagrass leaves as food source of *T. gratilla*.

The occurrence of some kind of foods in gut contents of sea urchin may indicate both, physical texture and chemical component in the seagrass leaves. Tertschnig (1989), explain that the preference for some type of seagrass was caused from some factors such as low tannin content (Chiroric acid). The high tannin content forced sea urchin *Tripneustes ventricosus* to decrease grazing seagrass leaves. Short and smooth texture leaves of *Th. hemprichii* were most common reasons for *T. gratilla* to select these kinds of leaves to graze (Kasim, 1999).

According to experiment by Mukai and Nojima (1985) grazing rates of *T. gratilla* in Papua New Guinean seagrass beds ranged from 2.0 to 5.7 g WW ind<sup>-1</sup> day<sup>-1</sup> (mean 3.6 g WW ind<sup>-1</sup> day<sup>-1</sup>) in single-animal experiment and 0.4 to 10.8 g WW ind<sup>-1</sup> day<sup>-1</sup> (mean 4.3 g WW ind<sup>-1</sup> day<sup>-1</sup>) in two-animal experiment, *T. gratilla* in Buton Island seagrass beds has biggest grazing rates around two times. In

Bolinao seagrass beds, Philippines, by field and laboratory measurement of *T. gratilla* and *Salmacis sphaeroides* consumption of seagrass, both sea urchins grazed more than 60 % of seagrasses, and total consumption was estimated 240 to 400 g DW m<sup>2</sup> day<sup>-1</sup> (Klumpp, *et al.*, 1993). Other research about grazing rates of sea urchin on seagrasses briefly were shown in **Table 3**.

Grazing activity of sea urchin on seagrasses has been a unique phenomenon correspondent directly with the biological and physical parameter. Food preference of sea urchin is caused by the availability of some kind of food which dominantly occurs around them. In *Thalassodendron ciliatum* meadows, Mombasa lagoon, *T. gratilla* has increased a high number of *T. ciliatum* dead shoot (Alcoverro and Mariani, 2002). In Buton Island seagrass beds, *Th. hemprichii* and *E. acoroides* are potential food sources of *T. gratilla*. All the seagrasses that put inside the cage from the grazing rate experiment were grazed by *T. gratilla*. In fact, immediately upon placing the seagrasses leaves on the substrat in the cages, *T. gratilla* began climbing the shoots and direct graze the leaves. The grazing rates on the seagrass leaves was related to different sized in diameter. During our experiment, the biggest sea urchins were grazing more seagrass leaves than the small sea urchins. Moor and McPherson (1965) reported that grazing rates of sea urchin is influenced by water current velocity, other condition such as food species (Vadas, 1968; Fuji 1967; Lowe 1974), Temperature (Agatsuma, *et al.*, 2000), season (Parcy, 1971; Moor and McPherson 1965), light intensity (Fuji, 1967), body size (Fuji, 1967; Leighton, 1968; Himmelman, 1969), and density (Greenway, 1976). The grazing time of *T. gratilla* is also influenced by the fluctuation of tidal range. The sea urchin hide under shoots of seagrasses during high tides when the current speed was high. It is considered that the most active graze of sea urchin is correlated to the low rate of flow.

The flow rate also affects to grazing rates of sea urchin.

## ACKNOWLEDGMENTS

I wish to thanks to Prof. Nawangsari Sugiri, Prof. H. Muhammad Eidman (Alm.), and Prof. Dr. H. Ridwan Affandi, for their constructive ideas and correction of this field research. Thanks for La Ode Abdul Umardin, Fedy As Ad, Asraf Mursalin and Mahyuddin for their field assistance. Special thanks for Erty M. Hatma and Cahyani Fitri Melania for their time and informal support. We also thank M. Kasim Naziri, Hj. Kasiah, Abdul Manan Hatma, Ny. Muriha, Mamnun, Danial and Indah Arfiyanti for their logistic support. Thanks to the laboratory of biology Haluoleo University for providing the laboratory facilities.

## REFERENCES

- Agatsuma, Y., Nakata, A. and Matsuyama, K. 2000. Seasonal foraging activity of the sea urchin *Strongylocentrotus nudus* on coralline flat in Oshoro Bay in southwestern Hokkaido, Japan. *Fisheries Science* 66 : 198-203.
- Alcoverro, T. and Mariani, S. 2002. Effects of sea urchin grazing on seagrass (*Thalassodendron ciliatum*) beds of a Kenyan lagoon. *Marine Ecological Progress Series* 226 : 255-263.
- Aziz, A. 1994. Food and feeding habit of sea urchin. *Oceana P30 LIPI XII*. 4 : 91 – 100. (In Indonesian)
- Bach, S.D. 1979. Standing crop, growth, and production of *Calcareaous siphonales* (Chlorophyta) in a south Florida lagoon. *Bulletin of Marine Science* 29 : 33 – 40.

- Bak, H.P. and Nojima, S. 1980. Immigration of a tropical sea urchin. *Astropiga radiata* (Laske) in a temperate eelgrass *Zostera marina* L. Patch; its feeding habit and grazing effect of the patch. *Publications of Amakusa Marine Biological Laboratory* 5 : 153 –169.
- Bouchon, C. 1980. Quantitative study of the scleractinian coral community of the Jordanian coast (Gulf of Aqaba, Red Sea); Preliminary results. *Tethys* 9 : 243 – 246.
- Cambridge, M.L., Chiffings, A.W., Brittan C., Moore L. and McComb A.J. 1986. The loss of seagrass in Cockburn Sound, Western Australia. 2 . *Aqua. Bot* 24 : 269 – 285.
- Camp, D.K., Cobb S.P. and Van Breedfield J.V. 1973. Overgrazing of seagrasses by a regular urchin, *Lytechinus variegates*. *Biol. Sci.* 23 : 37 – 38.
- Darsono, P. dan Sukarno. 1993. Biological aspect of sea urchin *Tripneustes gratilla* on Seagrass in Nusa Dua Bali. *Ocean of Indonesia* 26 : 13 – 25. (In Indonesian)
- den Hartog, C. 1970. *The seagrass of the world*. North Holland Publishing: Amsterdam. 275pp.
- Eklöfa, S. de la Torre-Castroa, M., Gullströmb, M., Ukuc, J., Muthigad, N., Lyimoe, T. and Bandeiraf, S.O., 2008, Sea urchin Overgrazing of Seagrasses: A review of Current Knowledge on Causes, Consequences, and Management, *Estuarine, Coastal and Shelf Science*, 79, Issue 4: 569-580
- Fuji, A. 1967. Ecological studies on the growth and food consumption of Japanese common littoral sea urchin, *Strongylocentrotus intermedius* (A. Agassiz). *Publication of Members of Faculty of Fisheries, Hokkaido University* 15 : 83-160.
- Greenway, M. 1976. The grazing of *Thalassia testudinum* in Kingston Harbour. Jamaica. *Aqua Bot* 2 : 117-126.
- Greenway, M. 1995. Tropic relationship of macrofauna within a Jamaica seagrass meadows and the role of the echinoid *Lytechinus variegatus* (Lamarck). *Bull. Mar. Sci* 56 : 719 – 736.
- Hattori, A., Aioi, K., Iizumi, H., Koike, I., Mukai, H., Nishihara, M., Nojima, S. and Yokohama, Y. 1985. *Studies on dynamic of the biological community in the tropical seagrass ecosystem in Papua New Guinean*. Report of the study support by the Grant-in-Aid of Ministry of Education. Culture and Science Japan. 49pp.
- Herring, P.J. 1972. Observation on the distribution and feeding habits of some littoral echinoids from Zanzibar. *J. Nat. His.* 6 : 169 – 175.
- Himmelman, J.H. 1969. Some aspect of the ecology *Strongylocentrotus drobachiensis* in eastern Newfoundland. MSc thesis. Memorial University. Newfoundland, 153pp.
- Huling, N.C. and Kirkman H. 1982. Further observation and data on seagrass along the Jordanian and Saudi Arabian coast of the gulf of Aqaba. *Tethys* 10 : 201 – 220.
- Ivlev, V.S. 1961. *Experimental ecology of the feeding of fishes*. Yale University Press, New Haven. 302 pp.
- Jafari, R.D. and Mahasneh R.M. 1984. The effect of seagrass grazing on the sexual maturity of the sea urchin *Tripneustes*

- gratilla* in the Gulf of Aqaba (Jordan).  
*Dirasat* 14 : 127 – 136.
- Kasim, M. 1999. Grazing activity and growth of sea urchin (*Tripneustes gratilla* L.) in Bone-Bone seagrass bed, Buton Island, Southeast Sulawesi, Indonesia. MSc. Thesis. Bogor Agricultural University. Indonesia. pp 38. (In Indonesian).
- Keller, B.D. 1983. Coexistence of sea urchin in seagrass meadows : an experimental analysis of competition and predation. *Eco.* 64 : 1581 – 1598.
- Kirkman, H. and Young, P.C. 1981. Measurement of health, and echinoderm grazing on *Posidonia oceanica* (L.) Delile. *Aqua. Bot.* 10 : 329 – 338.
- Klumpp, D.W., Salita-Espinosa, J.T. and Fortes, M.T. 1993. Feeding ecology and trophic role of sea urchin in tropical seagrass community. *Aqua. Bot.* 45 : 205 – 229.
- Larkum, A.W.D. and West, R.J. 1990. Long term changes of seagrass meadows in Botany Bay, Australia. *Aquatic Botany* 37 : 55 – 70.
- Lawrence, J.M. 1975. On the relationship between marine plant and sea urchin. *Oceanography Marine Biology Annual Review* 13 : 213 – 286.
- Leighton, D.L. 1968. A comparative study of food selection and nutrition in the abalone *Haliotis rufescens* Swainson, and the sea urchin, *Strongylocentrotus purpuratus* (Stimpson). PhD. Dissertation. Univ. California. San Diego. 197pp.
- Lipkin, Y. 1979. Quantitative aspect of seagrass communities, Particularly of those dominated by *Halophila stipulacea*, in Sinai (Northern Red Sea). *Aqua. Bot.* 7 : 119 – 128.
- Lowe, E.F. 1974. Absorption efficiencies, feeding rates and food preferences of *Lytechinus variegatus* (Echinodermata: Echinoidea) for select marine plants. MSc Thesis. University of South Florida. Tampa. 97pp.
- Moore, H.B. and McPherson, B.F. 1965. A contribution of the study of the productivity of the sea urchin *Tripneustes esculentus* and *Lytechinus variegatus*. *Bull. Mar. Sci.* 15 : 855-871.
- Mukai, H. and Nojima, S. 1985. A preliminary study on grazing and devication rates of seagrass grazer *Tripneustes gratilla*. (Echinodermata : Echinoids) in a Papua New Guinean seagrass bed. *Special Publication of Mukaishima Marine Biological Station* 1985 :185 – 192.
- Mukai, H., Nishihira, M. and Nojima, S. 1987. Distribution and biomass of predominant benthic animals study on dynamic of the biological community in tropical seagrass ecosystem in Papua New Guinea. The second report. Ocean Research Institute. University of Tokyo. 63 – 75.
- Oliveira, E. C., Corbisier, T.N., de Eston, V. R., Ambrosio Jr, O. 1997. Phenology of a seagrass (*Halodule wrightii*) bed on the southeast coast of Brazil. *Aqua. Bot.* 56 : 25 – 33.
- Parcy, J.A. 1971. Thermal acclimatization and acclimation in the echinoid, *Strongylocentrotus drobachiensis*. (O.F. Muller, 1776). PhD Dissertation. Memorial University of Newfoundland. 285pp.

- Tertschnig, W.P. 1989. Diel activity pattern and foraging dynamics of the sea urchin *Tripneustes ventricosus* in a tropical seagrass community and a reef environment (Virgin Island). *PSZN I: Mar. Ecol.* 10 : 3 – 21.
- Vadas, R. L. 1968. The ecology of *Agarum* and the kelp bed community. Ph D Dissertation. University of Washington. Washington. 280pp.
- Vadas, R. L., Fenchel, T. and Ogden, J.C. 1982. Ecological studies on the sea urchin *Lytechinus variegatus* and the algal-seagrass communities of the Miskito Cays, Nicaragua. *Aqua. Bot.* 14 : 139 – 150.
- Vaitilingon, D, Rasolofonirina, R and Jangoux, M. 2003, Feeding preferences, seasonal gut repletion indices, and diel feeding patterns of the sea urchin *Tripneustes graffilla* (Echinodermata: Echinoidea) on a coastal habitat off Toliara (Madagascar), *Marine biology*, 143 : 451-458.
- Zieman, J.C., Iverson, R.L. and Ogden, J.C. 1984. Herbivory effects on *Thalassia testudinum* leaf growth and nitrogen content. *Marine Ecology Progress Series* 15: 151 – 158.